



# 17<sup>th</sup> International Congress of the Mediterranean Phytopathological Union

New phytopathology frontiers of research  
and education in plant health and food safety

## BOOK OF ABSTRACTS



CIHEAM  
BARI

Italy

July 6-10, 2025

**P41**

**Evaluation of selected fungal biological control agents for the protection of grapevine pruning wounds against *Diplodia seriata***

A. FLOUDAS<sup>1</sup>, S. TESTEMPASIS<sup>1,2</sup>, A. FLARI<sup>1</sup>, E. DIMOU<sup>1</sup>, A. ELEFTHERIADOU<sup>1</sup>, G.S. KARAOGLANIDIS<sup>1</sup>

<sup>1</sup>Aristotle University of Thessaloniki, Faculty of Agriculture, Forestry and Natural Environment, Plant Pathology Laboratory, Thessaloniki, Greece. <sup>2</sup>Department of Agriculture, School of Agricultural Sciences, University of Western Macedonia, 53100 Florina, Greece. E-MAIL: gkarao@agro.auth.gr

Grapevine Trunk Diseases (GTDs) are among the most destructive grapevine diseases, causing significant economic losses due to vineyard decline and replanting costs. The lack of effective chemical measures highlights the urgent need for environmentally and consumer-friendly alternatives. Biological Control Agents (BCAs) offer a promising, sustainable approach, targeting multiple GTD-associated fungi. This field study was conducted in the viticultural region of Nemea to evaluate the efficacy of four BCAs - *Trichoderma atroviride*, *T. citrinoviride*, *T. ghanense*, *Talaromyces pinophilus*—and a mixture (*T. ghanense* + *T. pinophilus*) in controlling infections by *Diplodia seriata*, a key GTD pathogen in Greece. Two Greek grapevine cultivars with contrasting susceptibility - Roditis (susceptible) and Limnio (tolerant) were used. BCAs were applied as conidial suspensions on fresh pruning wounds, followed 24 hours later by artificial inoculation with *D. seriata*. Wounds were covered for two days to maintain moisture and promote infection. After six months, samples were collected and analyzed using a culture-based method. Pathogen reisolation was conducted on PDA medium, and all obtained fungal isolates were identified molecularly. The results revealed that *T. pinophilus* and its mixture with *T. ghanense* provided strong protection in the susceptible cultivar Roditis, substantially reducing pathogen recovery. Conversely, *T. atroviridae* was more effective in the tolerant cultivar Limnio, indicating a potential cultivar-specific interaction. All BCAs demonstrated the ability to colonize and persist on pruning wounds. These findings reinforce the potential of BCAs as effective, sustainable tools for GTD management and support their integration into future vineyard protection strategies.

This research was funded by the European project SHIELD4GRAPE (HORIZON-CL6-2023-BIODIV-01, NUMBER 101135088).